

IAP20 Rec'd PCT/PTO 06 FEB 2006

Advantageous combinations for inhalation of Nacystelyn and Bronchodilators**ABSTRACT**

- 5 The present invention provides with advantageous combinations for inhalation containing (A) L-Lysine-N-acetylcysteinate and (B) a bronchodilator agent and possibly pharmaceutical excipients. The composition can be formulated as a nebulizer, a metered dose inhaler or a dry powder inhaler.

10 BACKGROUND OF THE INVENTION

The main chronic respiratory diseases are Chronic obstructive pulmonary disease (COPD), asthma and Cystic Fibrosis (CF)

- 15 The definition of COPD originating from the Merck Manual of Diagnosis and Therapy, published by Merck Research Laboratories, Division of Merck & Co., Inc., Whitehouse Station, N.J., seventeen edition, 1999 is the following:

A disease characterized by chronic bronchitis or emphysema and airflow obstruction that is generally progressive, may be accompanied by airway
20 hyperreactivity, and may be partially reversible.

On the other hand, asthma is characterized by airway inflammation that is manifested by airway hyperresponsiveness to a variety of stimuli and by airway obstruction that is reversible spontaneously or in response to treatment;
25 reversibility may be incomplete in some patients.

Cystic fibrosis is another chronic inflammatory disease defined as follows :An inherited disease of the exocrine glands, primarily affecting the GI and respiratory systems, and usually characterized by COPD, exocrine pancreatic insufficiency,
30 and abnormally high sweat electrolytes.

BEST AVAILABLE COPY

The various terms used to quantify the pulmonary functions are described hereinbelow (Quanjer and al. , Eur. Respir. J., 1993, 6, Suppl. 16, 5-40, Official Statement of the European Respiratory Society)

- 5 Forced Vital Capacity (FVC) : is the volume of gas delivered during an expiration made as forcefully and completely as possible starting from full inspiration
- Forced expiratory volume in one second (FEV1) : is the volume of gas exhaled in a one second from the start of the forced vital capacity manoeuvre
- Peak Expiratory Flow Rate(PEFr) : is the maximal flow during a forced
- 10 expiratory vital capacity manoeuvre starting from a position of full inspiration

Chronic obstructive pulmonary disease (COPD), which includes chronic bronchiti and emphysema, is steadily increasing in frequency, possibly due to continued smoking, increasing air pollution, and the continued ageing of the population.

- 15 Prevalence: in the Global Burden of Disease Study conducted under the auspices of the WHO and the World Bank. The world-wide prevalence of COPD in 1990 was estimated to be 9.34/1,000 in men and 7.33/1,000 in women. However, these estimates include all ages and underestimate the true prevalence of COPD in adults. The prevalence of COPD is highest in countries where cigarette smoking has been,
- 20 or still is, very common, while the prevalence is lowest in countries where smoking is less common, or total tobacco consumption per individual is low.

Morbidity: the limited data that are available indicate that morbidity due to COPD increases with age and is greater in men than women, COPD is responsible for a significant part of physician visits, emergency department visits and

25 hospitalizations.

Mortality: COPD is currently the fourth leading cause of death in the world, and further increases in the prevalence and mortality of the disease can be predicted in the coming decades. In the US, COPD death rates are very low among people under age 45 but then increase with age, and COPD becomes the fourth or fifth

30 leading cause of death among those over 45.

COPD is characterized by edema (oedema) of the mucous membrane, which lines the anterior walls of the tracheobronchial tree. When the mucosa accumulates an

abnormal quantity of liquid, the profuse and thickened serous fluid excreted may seriously affect ventilation in the alveoli. The mucus resists movement up the walls of the tracheobronchial tree, normally efficiently accomplished by the cilia throughout the airways. Consequently, the serous fluid can form mucus plugs,
5 which can shut off alveoli or an entire airway depriving whole sections of the lung of oxygen-rich air.

Plugs of mucus in the tracheobronchial tree may only partially block the flow of air through the bronchioles. This partial blockage can create a turbulent flow of air, which forms bubbles on the surface of mucosa. When there are enough bubbles,
10 they become foam, which can clog airways and dramatically diminish respiration of the capillaries of the lungs.

The obstruction of the bronchi and bronchioles found in COPD is often a severely disabling condition. A wide variety of compounds including oral methylxanthines, oral and inhaled beta-adrenergic agonists, inhaled cromolyn sodium, inhaled
15 anticholinergics, and oral and inhaled corticosteroids have been tested. Despite the existence of these therapeutic tools, a large number of patients are not responsive to these medications or become non-responsive after a prolonged period of treatment. COPD is always accompanied by an important oxidative stress resulting from an oxidant/antioxidant imbalance, an excess of oxidants and/or a depletion of
20 antioxidants. Oxidative stress is thought to play an important role in the pathogenesis of a number of lung diseases, not only through direct injurious effects, but by involvement in the molecular mechanisms that control lung inflammation. A number of studies have shown an increased oxidant burden and consequently increased markers of oxidative stress in the airspaces, breath, blood,
25 and urine in smokers and in patients with COPD. The presence of oxidative stress has important consequences for the pathogenesis of COPD. These include oxidative inactivation of antiproteinases, airspace epithelial injury, increased sequestration of neutrophils in the pulmonary microvasculature, and gene expression of proinflammatory mediators. With regard to the latter, oxidative
30 stress has a role in enhancing the inflammation that occurs in smokers and patients with COPD, through the activation of redox-sensitive transcription factors such as

nuclear factor- κ B and activator protein-1, which regulate the genes for proinflammatory mediators and protective antioxidant gene expression.

5 Asthma is the most common form of bronchoconstrictive disease, which is completely different from COPD. Pathologically, asthma involves constriction of the bronchioles, hypertrophy of the muscles of the bronchioles, and a characteristic infiltrate of eosinophils.

Asthma is the third leading cause of preventable hospitalization in United States.
10 There are about 470.000 hospitalizations and more than 5.000 deaths a year from asthma. Asthma causes recurring episodes of coughing, wheezing, chest tightness, and difficult breathing. Asthma attacks can be life threatening. They can be prevented. Asthma is a chronic inflammatory disorder of the airways. Chronically inflamed airways are hyperresponsive; they become obstructed and airflow is
15 limited (by bronchoconstriction, mucus plugs, and increased inflammation) when airways are exposed to various stimuli, or triggers.

Common asthma triggers (that is, factors that make asthma worse) include viral infections; allergens such as domestic dust mites (in bedding, carpets, and fabric-upholstered furnishings), animals with fur, cockroach, pollens, and molds; tobacco
20 smoke; air pollution, exercise; strong emotional expressions; chemical irritants, and drugs (aspirin and beta blockers).

Asthma attacks (or exacerbations) are episodic, but airway inflammation is chronically present. Asthma is a chronic disorder requiring long-term management. For many patients, this means taking preventive medication every day.

25 Asthma can change over time. Asthma can be mild, moderate or severe; asthma attacks can be life-threatening. The severity of asthma varies among individuals, and it can change in one individual over time. Treatment decisions are made based on the severity of asthma.

30 In conclusion, it is clear that all chronic inflammatory diseases of the lungs like COPD and asthma present simultaneously important inflammation and / or

oxidation phenomenons of the lung tissue and a significant bronchoconstriction, making the breathing of patients very difficult. No efficient pharmaceutical treatment is currently available to threat all those symptoms simultaneously

- 5 A number of patents relating to the treatment of COPD or other pathologies of the respiratory tract have already been granted

The US Patent 6,153,187 describes a method of managing a patient having an accumulation of mucoid, mucopurulent or purulent material containing
10 glycosaminoglycans, the method comprising the step of administering at least one glycosaminoglycon degrading enzyme to the patient in an amount therapeutically effective to reduce at least one of the following: the visco-elasticity of the material, pathogens infectivity and inflammation.

- 15 The US Patent 5,969,421 describes a method for prevention of cell death, and method comprising simultaneously topically applying ACC and levulose as an application in an amount effective to result in a synergistic action protecting cells, and presenting cell damage caused by various agents. A pharmaceutical preparation including a combination of ACC and levulose.

20

The WO Patent 9635452 describes a pharmaceutical composition useful in the treatment of respiratory tract disorders. The composition comprises as active ingredients : (a) acetylcysteine, carbocysteine, erdocysteine or a pharmaceutically acceptable salt of any of theses; and (b) a beta 2 agonist, e.g. salbutamol,
25 terbutaline; and (c) an expectorant e.g. guaiphenesin, sodium citrate, ammonium chloride.

The WO Patent 0010598 describes a method of treating mucus hypersecretion, the causative factor in COPD, asthma and other clinical conditions involving COPD,
30 comprises administering a compound that inhibits exocytosis in mucus secreting cells or neurons that control or direct mucus secretion.

Some inventions describe the combination of a β_2 mimetics with an inhaled corticosteroid for the treatment of asthma and COPD. Examples of such inventions are EP416950, which describes the combination of salmeterol with beclomethasone; EP416951 relating to the combination of salmeterol with fluticasone or US 5,674,860 which relates to the combinations of formoterol and budesonide.

L-Lysine N-Acetylcysteinate (also called Nacystelyn or NAL) (US patent 4,847,282) is an active ingredient efficient in the treatment of chronic inflammatory diseases such as, but not limited to cystic fibrosis (CF) and chronic obstructive pulmonary disease (COPD) (Am. J. Respir. 2000, 161(3) – A72, pediatric pulmonology 22, 161-166 (1996), Free Rad. Biol. Med. 2002, 32(6), 492-502) NAL is actually a water-soluble salt of acetylcysteine (ACC), a widely used mucolytic. A number of experiments have demonstrated that NAL presents several advantages over ACC. The most important of them being the almost neutral pH of NAL ($\text{pH} \pm 6.5$) what makes its administration possible via the inhalation route without observing any bronchospasms, contrary to ACC, which presents an acidic pH ($\text{pH} \pm 2.2$) that can be responsible for reflex bronchospasms. This property allows the safe administration of NAL to patients by inhalation via all the inhaled technologies i.e.: nebulization, metered dose inhalers (MDIs) or drug powder inhalers, without the need of any buffering agents (Eur.Respir. J., 1994, 7(1), 81-87.

L-lysine N-acetylcysteinate (CAS N°89344-48-9) has been patented several years ago (US Patent 4,847,282, "Mucolytic acetylcysteine salts"). The principle of the invention was the synthesis of a water soluble acetylcysteine salt, consisting of reaction products of acetylcysteine with at least one basic amino-acid, the latter being preferably selected from the group comprising arginine, lysine, histidine, ornithine and glycine. A number of experiments have been performed to demonstrate that NAL presents several advantages over ACC, its parent molecule: the most important advantage is that NAL has an almost neutral pH ($\text{pH} = 6.5$),

what makes its administration possible via the inhalation route without observing any side effects (bronchospasms), contrary to ACC which is acidic (pH 2.2).

This allows the safe administration of NAL by inhalation by every galenical form available for this route e.g. nebulization, Metered Dose Inhalers (MDI) and Dry Powder Inhalers (DPI).

A number of experiments have been performed to assess the mucolytic, antioxidant and antiinflammatory properties of NAL in comparison to ACC. It is also important to mention that NAL is active at lower doses than ACC what makes its administration possible by inhalation via portable inhaler devices (DPI or MDI device).

This also greatly facilitates the administration for the patient and hence the compliance since there is no need to use a cumbersome nebulizer which requires an administration time of 10 to 30 minutes, what is difficult to accept for ambulatory patients like the majority of COPD or asthma patients. When NAL is formulated as a dry powder inhaler using a capsule device, the mean administration time for the patient is between 30 and 60 seconds.

The possibility of using a very convenient device and allowing to reach a high lung deposition of NAL is also very advantageous for the treatment of an ambulatory disease like COPD and asthma.

In the past, it has been demonstrated that surprisingly NAL presents a mucolytic and antioxidant activity superior to the activity of ACC (Vanderbist and al *Arzneim-Forschung/ Drug Research* (II) N°8, 1996, Nagy and al., *Pulmonary Pharmacology 2 Therapeutics* (1997) 10, 287-292, Gillissen and al *Respiratory Medicine* (1997), 91, 159-168, Tomkiewicz and al. *Pulmonary Pharmacology* (1995) 8, 259-265, Marriott and al, *Eur. Resp. J.*, 6 (suppl. 17), 438 (5) (abstract), 1993.

For some properties (mucolytic), the superiority of NAL may be explained while for other (antioxidant and antiinflammatory) effects, we can only make some hypothesis about the potential mode of action of the molecule.

Those mucolytic, antioxidant and antiinflammatory properties of NAL result in significant clinical improvement in patients. For instance, it has been demonstrated

in cystic fibrosis patients that the administration of NAL can significantly decrease the number and severity of lung exacerbations in comparison to placebo.

Bronchodilators are a very widely used treatment of chronic respiratory diseases. Although they can act through various mechanisms of action, they all result in the dilatation of the airways of the respiratory tract, resulting in an improvement of patient's lung functions such as Forced Vital Capacity (FVC), Forced Expiratory Volume in one second (FEV1) and Peak Expiratory Flow Rate (PEFR). There are two main classes of bronchodilators: the sympaticomimetics, including the short-acting and the long-acting β 2-mimetics, and the anticholinergics.

Short-action β 2-mimetics include (but are restricted to) salbutamol, terbutaline, fenoterol, pirbuterol or tulobuterol. Each active ingredient cited can be used as the base or as an acceptable pharmaceutical salt. The long-acting β 2 mimetics include (but are not restricted to) formoterol and salmeterol. Each molecule can be used as base or as an acceptable pharmaceutical salt. The anticholinergic drugs include (but are not restricted to) ipratropium, oxitropium or tiotropium.

In general, bronchodilators are used in asthma and chronic obstructive pulmonary diseases but can be administered in almost every inflammatory disease of the lung. Never described was a combination for inhalation of the mucoactive agent L-lysine N-acetylcysteinate (Nacystelyn or NAL) with a bronchodilator agent.

FIELD OF THE INVENTION

Chronic respiratory diseases involve complexes mechanisms and require the adminsitration of different kinds of treatment to patients among with antibiotics, corticosteroids, bronchodilators, mucolytics,.... The field of the present invention is to provide an advantageous , efficient and safe treatment of such diseases by administering an combination for inhalation comprising the mucoactive agent

Nacystelyn and at least one bronchodilator agent. Advantageous combinations are disclosed in the attached claims.

It has now suprisingly been found that NAL and bronchodilators can
5 advantageously be combined to be used by inhalation in the treatment of chronic
respiratory diseases such as cystic fibrosis, chronic obstructive pulmonary disease
or asthma. NAL has a duration of action that makes its administration possible
from 1 to 4 times a day. The bronchodilator class of molecules include molecules
10 with very long duration of action which have to be administered only once a day
(tiotropium) but also long acting β 2-mimetics which are usually administered twice
a day like formoterol and salmeterol. Finally there are short-acting bronchodilators
such as salbutamol, terbutaline, ipratropium or oxitropium which have to be
administered 4 to 6 times a day. Consequently, NAL can be combined in a single
15 pharmaceutical form with every type of bronchodilators. NAL and bronchodilators
offer complementary modes of action which make their administration particularly
profitable for the patients. Indeed, NAL, through its mucolytic activity may allow
to the bronchodilator to be deposited more deeply in the lungs after inhalation i.e.
in its target tissue, the bronchi and the bronchioli, so permitting a better efficacy of
the bronchodilator agent. On the other hand, through its action of dilatation of the
20 conducting airways, the bronchodilator agent will allow to NAL to also be
deposited deeply in the lungs so guarantying an optimal antiinflammatory and
antioxydant of NAL, all along the respiratory tract.

Using such combination therapy, medicaments which result in a significant
improvement in lung function may be prepared. In another aspect, using the
25 combination therapy of the invention, medicaments which provide improved
control of obstructive or inflammatory airways diseases, or a reduction in
exacerbation's of such diseases, may be prepared. In a further aspect, using
compositions of the invention, medicaments which can be used on demand in
rescue treatment of obstructive or inflammatory airways diseases, or which reduce
30 or eliminate the need for treatment with short-acting rescue medicaments such as
salbutamol or terbutaline, may be prepared; thus medicaments based on

compositions of the invention facilitate the treatment of an obstructive or inflammatory airways disease with a single medicament.”

The combination allowing an administration twice or once a day will be preferred to combinations requiring more administrations because they increase the comfort and the compliance of the patients. NAL and the bronchodilator can be
5 administered simultaneously, separately or sequentially. The combination of NAL and the bronchodilator agent can be formulated as a nebulizer, as metered dose inhalers (MDI) or as a drug powder inhaler (DPI).

For a more easier understanding of which kind of combination the present
10 invention is relating, in the following of the present text, the letter (A) will be used to design Nacystelyn and the letter (B) will be used to design the bronchodilator agent. The present invention can for instance, consist in the administration of the medicament or pharmaceutical composition as hereinbefore described, i.e. with (A) and (B) in admixture or separate, by inhalation, i.e. (A) and (B) or the mixture
15 thereof are in inhalable form. The inhalable form of the medicament i.e. of (A) and (B) may be, for example, an atomizable composition such as an aerosol comprising the active ingredient, i.e. (A) and (B) separately or in admixture, in solution or dispersion in a propellant, or a nebulizable composition comprising a dispersion of the active ingredient in an aqueous, organic or aqueous/organic medium. For
20 example, the inhalable form (suitable for inhalation) of the medicament may be an aerosol comprising a mixture of (A) and (B) in solution or dispersion in a propellant, or a combination of an aerosol containing (A) in solution or dispersion in a propellant with an aerosol containing (B) in solution or dispersion in a propellant. In another example, the inhalable form is a nebulizable composition
25 comprising a dispersion of (A) and (B) in an aqueous, organic or aqueous/organic medium, or a combination of a dispersion of (A) in such a medium with a dispersion of (B) in such a medium.

An aerosol composition suitable for use as the inhalable form of the medicament may comprise the active ingredient in solution or dispersion in a propellant, which
30 may be chosen from any of the propellants known in the art. Suitable such propellants include hydrocarbons such as n-propane, n-butane or isobutane or mixtures of two or more such hydrocarbons, and halogen-substituted hydrocarbons,

for example fluorine-substituted methanes, ethanes, propanes, butanes, cyclopropanes or cyclobutanes, particularly 1,1,1,2-tetrafluoroethane (HFA134a) and 1,1,1,2,3,3,3-heptafluoropropane (HFA227), or mixtures of two or more such halogen-substituted hydrocarbons. Where the active ingredient is present in suspension in the propellant, i.e. where it is present in particulate form dispersed in the propellant, the aerosol composition may also contain a lubricant and a surfactant, which may be chosen from those lubricants and surfactants known in the art, like for instance sorbitan oleate and derivatives. Other suitable aerosol compositions include surfactant-free or substantially surfactant-free aerosol compositions. The aerosol composition may contain up to about 5% by weight, for example 0.002 to 5%, 0.01 to 3%, 0.015 to 2%, 0.1 to 2%, 0.5 to 2% or 0.5 to 1%, by weight of the active ingredients, based on the weight of the propellant. Where present, the lubricant and surfactant may be in an amount up to 5% and 0.5% respectively by weight of the aerosol composition. The aerosol composition may also contain a co-solvent such as ethanol in an amount up to 30% by weight of the composition, particularly for administration from a pressurized metered dose inhalation device.

In another embodiment of the invention, the inhalable form is a dry powder, i.e. (A) and (B) are present in a dry powder comprising finely divided (A) and (B) optionally together with a finely divided pharmaceutically acceptable carrier, which is preferably present and may be one or more materials known as pharmaceutically acceptable carriers, preferably chosen from materials known as carriers in dry powder inhalation compositions, for example saccharides, including monosaccharides, disaccharides, polysaccharides and sugar alcohols such as arabinose, glucose, fructose, ribose, mannose, sucrose, trehalose, lactose, maltose, starches, dextran or mannitol. An especially preferred carrier is lactose. The dry powder may be in capsules of hard gelatin or hydroxypropylmethylcellulose, or in blisters, for use in a dry powder inhalation device, preferably in dosage units of (A) and (B) together with the carrier in amounts to bring the total weight of powder per dose to from 5 mg to 50 mg. Alternatively, the dry powder may be contained as a reservoir in a multi-dose dry powder inhalation device.

In the finely divided particulate form of the medicament, and in the aerosol composition where the active ingredient is present in particulate form, the active ingredient may have an average particle diameter of up to about 10 μm , for example 0.1 to 5 μm , preferably 1 to 5 μm . The solid carrier, where present, generally has a maximum particle diameter up to 300 μm , preferably up to 212 μm , and conveniently has a mean particle diameter of 100 to 160 μm , e.g. 100 to 125 μm . The particle size of the active ingredient, and that of a solid carrier where present in dry powder compositions, can be reduced to the desired level by conventional methods, for example by grinding in an air-jet mill, ball mill or vibrator mill, microprecipitation, spray-drying, lyophilisation or recrystallisation from supercritical media. the carrier material can also consist in a mix of different materials in order to optimize the properties of the dry powder inhaler formulation. The inhalable medicament may be administered using an inhalation device suitable for the inhalable form, such devices being well known in the art. Accordingly, the invention also provides a pharmaceutical product comprising a medicament or pharmaceutical composition as hereinbefore described in inhalable form as hereinbefore described in association with one or more inhalation devices. In a further aspect, the invention provides an inhalation device, or a pack of two or more inhalation devices, containing a medicament or pharmaceutical composition as hereinbefore described in inhalable form as hereinbefore described.

Examples :

The invention is additionally illustrated in connection with the following examples, which are considered to be illustrative of the present invention. It should be understood, however, that the invention is not limited to the specific details of the Examples.

Example 1 : combination: NAL / Ipratropium 50 / 0.25 mg – powder for nebulization

Formula per vial :

Nacystelyn : 50 mg
ipratropium bromide : 0.25 mg
Mannitol : 44.7 mg
sodium edatate (EDTA): 0.10 mg

5 reconstitution is made with 2 ml of deionized water (degassed for 30 minutes)

Process : The combination was manufactured using a freeze-drying process. (freeze-drying machine : GT2 (leybold-Heraeus). The solution of Nacystelin and ipratropium bromide (in a solvent, advantageously in an aqueous solvent or water
10 containing solvent) with mannitol was filled (1 ml) into Typr I glass vials (Gaasch packaging) under constant nitrogen gas bubbling. The vials were gassed with nitrogen and partly closed with rubber stoppers (FM257/2, Helvoet Pharma) and placed inside the Freeze-dryer. After lyophilization, the chamber was vented with nitrogen gas and the vials were automatically closed and sealed with Al caps.

15

The so obtained powder is then suitable for ensuring a nebulization of the compounds with an aqueous medium.

Example 2 : NAL / Oxitropium bromide 4 / 0.1 mg / puff

20

Formula

ingredient	Amount / puff (mg / puff)
Nacystelyn	2.0
Oxitropium bromide	0.1
lecithin	7.5
monofluorotrichloroethane	25.0
difluorodichloromethane	60.0

Example 3 : NAL / salmeterol 3 / 0.050 mg / puff

The salmeterol used was under the form of salmeterol xinafoate. 73.2 mg of
 5 salmeterol xinafoate are equivalent to 50.0 mg of salmeterol base

Formula :

ingredient	Amount / puff (mg / puff)
Nacystelyn	4
salmeterol xinafoate	0.073
lecithin	7.5
monofluorotrichloroethane	20.0
difluorodichloromethan	50.0

10 **Example 4 : combination NAL / Formoterol - dry powder inhaler formulation**

Formula :

ingredient	Amount / dose(mg / dose)
Nacystelyn	4.0
Formoterol fumarate	0.006
Lactose monohydrate	5.90

15 **Process :** formoterol fumarate (with a weight average particle size comprised between 1 and 5 μm) is pre-blended with a small fraction of lactose (1/10 of the total amount of lactose with a weight particle size comprised between 100 and

300 μ m, such as about 160 μ m,) in a cubic mixer (Turbula). NAL (with a weight average particle size comprised between 1 and 5 μ m) is blended in a planetary mixer (Colette) with the remaining part of lactose (9/10 of the amount of lactose) for 10 minutes on speed 1. The pre-blend formoterol / lactose is then added to the
5 NAL / lactose mix in the planetary mixer (sandwich mix) and the final blend is mix at speed 1 for another 10 minutes. The resulting powder is then manually filled into Multidose DPI device (Mulidose MIAT Inhaler, Milano, Italy). Another part of the mix has been filled into size 3 hydroxypropylmethylcellulose capsules to be used with a single dose DPI device (Miat Monodose inhaler).

10

Example 5 : combination NAL / Formoterol fumarate - dry powder inhaler formulation 10 / 0.012 mg

Formula :

15

ingredient	Amount / capsule (mg / capsule)
Nacystelyn	10.0
Formoterol fumarate	0.012
Lactose monohydrate	8.0
Anhydrous lactose	22.0

The mixing process was the same as in example 4.

The powder was filled into transparent – transparent , size 3 hydroxypropylmethylcellulose capsules and administered with the monodose Miat
20 inhaler (Miat, Milano, Italy).

Example 6 : Stability data on NAL / formoterol DPI from example 5

Stability data on NAL / formoterol capsules packaged in aluminum / aluminum blisters

5

A) 25 °C / 60 % RH

	T0	T3 Months	T6 Months	T12 Months	T18 Months	T24 Months
<u>Assay</u>						
formoterol (%)	99.8	98.9	99.6	99.2	99.1	99.3
NAL (%)	101.2	100.6	100.4	100.7	100.8	100.2
<u>Impurities</u>						
a) formoterol						
impurity 52008RC01 (%)	0.2	0.3	0.2	0.3	0.4	0.5
total impurities (%)	0.2	0.3	0.3	0.4	0.5	0.6
b) NAL						
NN-diacecylcystine (%)	0.15	0.20	0.22	0.30	0.32	0.46
total impurities (%)	0.20	0.22	0.25	0.26	0.27	0.28
<u>Fine particle Dose</u>						
formoterol (µg)	3.7	3.6	3.8	3.6	3.9	3.8
NAL (mg)	3.2	3.1	3.1	3.0	3.2	3.1

The stability tests were carried out in order to establish the presence of impurities and the fine particle dose after preparation, as well as after a storage period of 3 months, 6 months, 12 months, 18 months and 24 months at 25°C with a relative humidity of 60%. This test shows the stability of the composition with respect to the formation of impurities, as well as with respect to the fine particle dose for the formoterol and for the NAL.

10

**Example 7 : combination NAL / Salmeterol - dry powder inhaler formulation
10 / 0.073 mg (equal to 50.0 mg of salmeterol base)**

ingredient	Amount / capsule (mg / capsule)
Nacystelyn	10.0
Salmeterol xinafoate	0.073
Lactose monohydrate	8.0
Anhydrous lactose	22.0

5

**Example 8 : Fine particle Dose (FPD) obtained with the DPI combination
NAL / Salmeterol of example 7 versus the marketed DPI form of salmeterol
DPI (Serevent 50 µg, Diskus)**

10

From an In vitro comparative deposition of salmeterol from (i) the combination
NAL / salmeterol 10 / 0.025 mg of the current invention versus (ii) the Serevent 50
µg, Diskus (Deposition measured with a Multistage Liquid Impinger, 4 liters of
air, n=3), it appears that the fine particle dose obtained with the combination of the
invention (NAL + 25µg salmeterol) was 10.9 µg salmeterol, while said fine
particle dose was 9.89 µg salmeterol for the Serevent 50µg salmeterol

20

CLAIMS

-
1. A pharmaceutical combination or composition for inhalation a fixed
5 combination (A) L-lysine N-acetylcysteinate and (B) a bronchodilator agent
for simultaneous, sequential or separate administration by inhalation in the
treatment of an inflammatory or obstructive respiratory disease.
 2. The composition or combination of claim 1, wherein the bronchodilator
agent belongs to the class of β 2-mimetics.
 - 10 3. The composition or combination of claim 1, wherein the bronchodilator
agent belongs to the class of long-acting β 2-mimetics.
 4. The composition or combination of claim 3, wherein the bronchodilator
agent is formoterol or a solvate thereof, or salmeterol or a salt thereof, or
mixtures thereof.
 - 15 5. The composition or combination of claim 2, wherein the bronchodilator
agent is a short-acting β 2-mimetic such as but not limited to salbutamol,
terbutaline, pirbuterol, fenoterol, tulobuterol, and mixtures thereof.
 6. The composition or combination of claim 1, wherein the bronchodilator
agent belong to the class of anticholinergics.
 - 20 7. The composition or combination of claim 6, wherein the bronchodilator
agent is tiotropium, oxitropium, ipratropium and mixtures thereof.
 8. The composition or combination of claim 1, wherein the weight ratio of (A)
to (B) is from 1:10 to 1:10 000, preferably from 1:100 to 1:2000, most
preferably from 1:200 to 1:1000.
 - 25 9. The composition or combination of claim 1, wherein the composition or
combination contained in addition of (A) and (B) at least one
pharmaceutically acceptable carrier.
 10. The composition or combination of claim 1 suitable for an inhalable dry
powder inhaler comprising micronized particles of (A) L-lysine N-
30 acetylcysteinate and (B) micronized particles of (B) a bronchodilator agent,
advantageously mixed with one or more acceptable excipients(s)

11. The composition or combination of claim 10, wherein the composition or combination is in a form suitable for inhalation through a monodose inhaler device.
12. The composition or combination of claim 10, wherein the composition or combination is in a form suitable for inhalation through a multiple-dose inhaler.
13. The composition or combination of claim 10, containing in addition to (A) and (B) at least one saccharide.
14. The composition or combination of claim 13, where the saccharide is lactose.
15. The composition or combination of claim 1, wherein the composition or combination is an inhalable pressurized metered dose inhaler.
16. The composition or combination of claim 15, wherein the composition or combination contains in addition to (A) and (B) at least a propellant wherein (A) and (B) are either dissolved or dispersed.
17. The composition or combination of claim 15, wherein the propellant is chlorofluorocarbon derivative.
18. The composition or combination of claim 15, where the propellant is a hydrofluorocarbon derivative.
19. The composition or combination of claim 1, wherein the composition or combination is an inhalable nebulizable composition or combination comprising a solution and/or a dispersion of (A) and (B) in an aqueous medium.
20. The composition or combination of claim 1, which is a dry powder inhaler in a capsule containing from 2 to 20 µg of L-lysine N-acetylcysteinate and from 2 to 20 µg of formoterol or a solvate thereof.
21. The composition or combination of claim 1, which is a dry powder inhaler in a capsule containing from 2 to 20 µg of L-lysine N-acetylcysteinate and from 10 to 100 µg of salmeterol or a solvate thereof.
22. The composition or combination of claim 1, which is a dry powder inhaler in a capsule containing from 2 to 20 µg of L-lysine N-acetylcysteinate and from 10 to 100 µg of tiotropium or a solvate thereof.

23. A method of treating an inflammatory or obstructive respiratory disease which comprises administering to a subject in need of such treatment effective amounts of (A), as defined in claim 1, and (B) as defined in claim 1.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/BE 03/00134

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/195 A61K31/167 A61K31/216 A61K31/138 A61P11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 48839 A (SEGAL CATHERINE A ;WARNER LAMBERT CO (US)) 5 November 1998 (1998-11-05)	
A	WO 03 035137 A (RESPIRICS INC) 1 May 2003 (2003-05-01)	
A	EP 0 876 814 A (PHARLYSE SA) 11 November 1998 (1998-11-11)	
A	US 4 847 282 A (DEBOECK ARTHUR M) 11 July 1989 (1989-07-11) cited in the application	

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

25 November 2003

Date of mailing of the international search report

03/12/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Stienon, P

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat^l publication No

PCT/BE 03/00134

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9848839	A	05-11-1998	AU 6878098 A BR 9809022 A CN 1253508 T EP 0979105 A1 JP 2001524108 T WO 9848839 A1	24-11-1998 01-08-2000 17-05-2000 16-02-2000 27-11-2001 05-11-1998
WO 03035137	A	01-05-2003	US 2003075172 A1 WO 03035137 A2	24-04-2003 01-05-2003
EP 0876814	A	11-11-1998	EP 0876814 A1 AU 7201298 A WO 9850015 A1 DE 69817774 D1 EP 0964675 A1	11-11-1998 27-11-1998 12-11-1998 09-10-2003 22-12-1999
US 4847282	A	11-07-1989	LU 84095 A1 AT 26109 T DE 3370477 D1 EP 0092287 A2	16-12-1983 15-04-1987 30-04-1987 26-10-1983

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference RP/PCT/03-16	FOR FURTHER ACTION <small>see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.</small>	
International application No. PCT/BE 03/00134	International filing date (day/month/year) 06/08/2003	(Earliest) Priority Date (day/month/year)
Applicant GALEPHAR M/F		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :

☐ contained in the international application in written form.

☐ filed together with the international application in computer readable form.

☐ furnished subsequently to this Authority in written form.

☐ furnished subsequently to this Authority in computer readable form.

☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No. _____

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☒ None of the figures.

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/BE 03/00134

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/195 A61K31/167 A61K31/216 A61K31/138 A61P11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 48839 A (SEGAL CATHERINE A ;WARNER LAMBERT CO (US)) 5 November 1998 (1998-11-05) ---	
A	WO 03 035137 A (RESPIRICS INC) 1 May 2003 (2003-05-01) ---	
A	EP 0 876 814 A (PHARLYSE SA) 11 November 1998 (1998-11-11) ---	
A	US 4 847 282 A (DEBOECK ARTHUR M) 11 July 1989 (1989-07-11) cited in the application -----	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

25 November 2003

Date of mailing of the international search report

03/12/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Stienon, P

THIS PAGE BLANK (USPTO)


INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/BE 03/00134

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9848839	A	05-11-1998	AU 6878098 A BR 9809022 A CN 1253508 T EP 0979105 A1 JP 2001524108 T WO 9848839 A1	24-11-1998 01-08-2000 17-05-2000 16-02-2000 27-11-2001 05-11-1998
WO 03035137	A	01-05-2003	US 2003075172 A1 WO 03035137 A2	24-04-2003 01-05-2003
EP 0876814	A	11-11-1998	EP 0876814 A1 AU 7201298 A WO 9850015 A1 DE 69817774 D1 EP 0964675 A1	11-11-1998 27-11-1998 12-11-1998 09-10-2003 22-12-1999
US 4847282	A	11-07-1989	LU 84095 A1 AT 26109 T DE 3370477 D1 EP 0092287 A2	16-12-1983 15-04-1987 30-04-1987 26-10-1983



THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)